



Office de la propriété  
intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada

10/523761

DT12 Rec'd PCT/PTO 07 FEB 2005

*Bureau canadien  
des brevets*  
Certification

*Canadian Patent  
Office*  
Certification

La présente atteste que les documents  
ci-joints, dont la liste figure ci-dessous,  
sont des copies authentiques des docu-  
ments déposés au Bureau des brevets.

This is to certify that the documents  
attached hereto and identified below are  
true copies of the documents on file in  
the Patent Office.

Specification as originally filed, with Application for Patent Serial No: **2,395,819**, on  
August 13 2002, by **BERNARD CHARLES SHERMAN**, for "Dual-Spike Release  
Formulation for Oral Drug Delivery".

  
Agent certificateur/Certifying Officer

February 1, 2005  
Date

Canada

(CIPO 68)  
31-03-04

OPIC  CIPO

**ABSTRACT**

5 A composition which achieves release of a drug in two spikes, and which  
comprises particles, wherein the particles consist of a homogenous mixture  
which comprises the drug and an enteric polymer.

10

15

20

25

30

**DUAL-SPIKE RELEASE FORMULATION**  
**FOR ORAL DRUG DELIVERY**

**Background of the Invention**

5

Methylphenidate is a compound used to treat hyperactivity and attention deficit disorder (ADD) in children.

10

Methylphenidate is sold in the United States and elsewhere under the trade name Ritalin™. More particularly, it is sold as Ritalin™ immediate-release tablets for oral administration containing methylphenidate as the hydrochloride salt in strengths of 5, 10 and 20 mg. The drug is also available in Ritalin SR™ tablets which contain methylphenidate as the hydrochloride salt in strength of 20 mg in a slow-release formulation.

15

Ritalin SR™ is not suitable for optimal treatment of ADD because it produces a continuous slow release, whereas methylphenidate requires a spike or "sawtooth" (i.e. multi-spike) profile to have maximum efficacy in the treatment of ADD. For this reason, ADD has usually been treated with prompt release tablets given two or three times daily.

20

It would thus be desirable to have a dosage form that could be administered as a single dose in the morning and would release the drug in two spikes at least one hour apart.

25

Such a product is now being introduced into the United States market under the tradename Ritalin LA™ capsules in strengths of 20, 30 and 40 mg. These capsules are made in accordance with the teachings of U.S. patent application number US2001/0046472. Each capsule contains two different types of beads. Half of the dose is in immediate-release beads, which release their drug content in the stomach. The other half of the dose is in delayed-

30

™ - Trademark

release beads. These are beads which are enteric coated; that is to say they are coated with a polymer that is insoluble at acidic gastric pH, but soluble at intestinal pH. Hence, the delayed-release beads release these drug contents only after the pellets reach the small intestine and the enteric coating dissolves.

While the formulation of Ritalin LA capsules achieves the desired result of two spikes, the formulation is complex and expensive to manufacture. It requires production of two different types of beads, and also requires capsule-filling equipment that is capable of filling two different types of beads into a single capsule.

In view of this prior art, the objective of the present invention is to enable a formulation that will release the drug content in two peaks from a single type of bead, pellet or granule.

#### Description of the Invention

It has been found that particles in the form of beads, pellets or granules can be made of a single type that release a drug in two spikes, the first spike occurring promptly upon the particles reaching the stomach, and the second peak occurring after the particles have traveled into the small intestine.

The present invention is a pharmaceutical composition comprising such particles.

The particles consist of an essentially homogenous mixture, which comprises both a water-soluble drug and an enteric polymer. It has been found that if the particles are very small and the ratio of enteric polymer to the drug is low,

then all or essentially all of the drug will be promptly released in the stomach. However, as the size of the particles is increased and the ratio of enteric polymer to drug is increased, it is found that only the drug content that is  
5 closest to the surface of the particles leaches out of the mixture and dissolves in the acidic gastric fluid; dissolution then ceases, as the portion of the drug that is further into the particles is protected against dissolution by the remaining enteric polymer in the outer portion of the particle from which the drug has leached. Dissolution then essentially ceases until the particles reach  
10 the small intestine and the pH is high enough so that the enteric polymer dissolves to release the balance of the drug.

By trial and error, a ratio of the enteric polymer to drug and a particle size range may be found such that approximately half of the drug will be promptly  
15 released in the stomach and the other half will be released in the small intestine.

The ratio of enteric polymer to drug by weight will preferably be at least 2 and less than 100, will more preferably be from about 4 to about 50 and will most  
20 preferably be from about 10 to about 20.

The drug may be methylphenidate or a salt thereof, preferably methylphenidate hydrochloride.

25 The enteric polymer may be any enteric polymer acceptable for use in pharmaceuticals such as, for example, polyvinyl acetate phthalate, hydroxypropyl methylcellulose phthalate, cellulose acetate phthalate, cellulose acetate trimethitate, hydroxypropyl methylcellulose acetate succinate, and methacrylic acid copolymer type C. Most preferred is polyvinyl acetate  
30 phthalate.

The invention will be better understood from the following example, which is intended to be illustrative and not intended to limit the scope of the invention.

5 Drug and enteric polymer were mixed in the following proportions.

Methylphenidate Hydrochloride	20 parts
Polyvinyl acetate phthalate	<u>260 parts</u>
	280 parts

10

The mixture was compressed into slugs (large tablets) on a tablet press. The slugs were then ground up through #8 screen (8 wires to the inch). The resulting granules were then shaken on a #16 screen, and the portion that went through the #16 screen was discarded. The granules that remained on  
15 the screen were then filled into capsules at a net fill of 280 mg per capsule, so that each capsule contained granules comprising 20 mg of methylphenidate hcl.

Dissolution of the capsules was then tested in United States Pharmacopoeia  
20 apparatus #2 at 50 RPM, in both simulated gastric fluid and simulated intestinal fluid of pH6.8. It was found that, in simulated gastric fluid, approximately half of the drug was released promptly (within 30 minutes) and dissolution then ceased.

On the other hand, in simulated intestinal fluid, dissolution was essentially  
25 complete within 2 hours. It follows that when these capsules are ingested, approximately half of the drug content will be released promptly in the stomach, and the rest will be released within about 2 hours after the particles reach the more alkaline fluid of the small intestine.

**Claims**

- 5 1. A composition for the oral administration of a drug which achieves drug release in two spikes and which comprises particles, wherein the particles consist of a homogenous mixture which comprises the drug and an enteric polymer.
- 10 2. A composition of claim 1 wherein the ratio of enteric polymer to drug is at least 2 and less than 100.
3. A composition of claim 1 wherein the ratio of enteric polymer to drug is from about 4 to about 50.
- 15 4. A composition of claim 1 wherein the ratio of enteric polymer to drug is from about 10 to about 20.
- 20 5. A composition of any of claims 1 to 4 wherein the enteric polymer is polyvinyl acetate phthalate.
6. A composition of any of claims 1 to 5 wherein the drug is methylphenidate or a salt thereof.

25

30